

### Clinical Pharmacy Program Guidelines for Repatha

Program	Prior Authorization
Medication	Repatha (evolocumab)
Issue Date	5/2015
Pharmacy and Therapeutics Approval Date	11/2017
Effective Date	1/2018

#### 1. Background:

Repatha™ (evolocumab) is a PCSK9 (proprotein convertase subtilisin kexin type 9) inhibitor antibody indicated as an adjunct to diet and:

- Maximally tolerated statin therapy for treatment of adults with heterozygous familial hypercholesterolemia (HeFH) or clinical atherosclerotic cardiovascular disease (CVD), who require additional lowering of low density lipoprotein cholesterol (LDL-C)
- Other LDL-lowering therapies (e.g., statins, ezetimibe, LDL apheresis) in patients with homozygous familial hypercholesterolemia (HoFH) who require additional lowering of LDL-C

#### 2. Coverage Criteria:

##### A. Primary Hyperlipidemia

##### 1. Initial Therapy

a. **Repatha** will be approved based on **all** of the following criteria:

(1) **One** of the following diagnoses:

(a) Heterozygous familial hypercholesterolemia (HeFH) as confirmed by **one** of the following\*:

i. **Both** of the following:

1. Pre-treatment LDL-C greater than 190 mg/dL (greater than 155 mg/dL if less than 16 years of age)

-AND-

2. **One** of the following:

Confidential and Proprietary, © 2017 UnitedHealthcare Services Inc.

- a. Family history of myocardial infarction in first-degree relative < 60 years of age
- b. Family history of myocardial infarction in second-degree relative < 50 years of age
- c. Family history of LDL-C greater than 190 mg/dL in first- or second-degree relative
- d. Family history of familial hypercholesterolemia in first- or second-degree relative
- e. Family history of tendinous xanthomata and/or arcus cornealis in first- or second degree relative

**-OR-**

ii. **Both** of the following:

1. Pre-treatment LDL-C greater than 190 mg/dL (greater than 155 mg/dL if less than 16 years of age)

**-AND-**

2. Submission of medical records (e.g., chart notes, laboratory values) documenting **one** of the following:

- a. Functional mutation in LDL, apoB, or PCSK9 gene\*
- b. Tendinous xanthomata
- c. Arcus cornealis before age 45

**-OR-**

(b) Atherosclerotic cardiovascular disease (ASCVD) as confirmed by one of the following:

- i. Acute coronary syndromes
- ii. History of myocardial infarction
- iii. Stable or unstable angina
- iv. Coronary or other arterial revascularization
- v. Stroke
- vi. Transient ischemic attack
- vii. Peripheral arterial disease presumed to be of atherosclerotic origin

**-AND-**

(2) Submission of medical records (e.g., chart notes, laboratory values) documenting **one** of the following [prescription claims history may be

used in conjunction as documentation of medication use, dose, and duration]:

- (a) Patient has been receiving at least 12 consecutive weeks of **high-intensity statin therapy** [i.e. atorvastatin 40-80 mg, rosuvastatin 20-40 mg] and will continue to receive high-intensity statin at maximally tolerated dose

**-OR-**

- (b) **Both** of the following:

- i. Patient is unable to tolerate **high-intensity statin** as evidenced by **one** of the following intolerable and persistent (i.e. more than 2 weeks) symptoms:
- Myalgia (muscle symptoms without CK elevations)
  - Myositis (muscle symptoms with CK elevations < 10 times upper limit of normal [ULN])

**-AND-**

- ii. **One** of the following:

a. Patient has been receiving at least 12 consecutive weeks of **moderate-intensity statin** [i.e. atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin  $\geq$  20 mg, pravastatin  $\geq$  40 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily or Livalo (pitavastatin)  $\geq$  2 mg] and will continue to receive a moderate-intensity statin at maximally tolerated dose

**-OR-**

b. Patient has been receiving at least 12 consecutive weeks of **low-intensity statin** [i.e. simvastatin 10 mg, pravastatin 10-20 mg, lovastatin 20 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1 mg] therapy and will continue to receive a low-intensity statin at maximally tolerated dose

**-OR-**

- (c) Patient is unable to tolerate **low or moderate-, and high-intensity statins** as evidenced by **one** of the following:

- i. **One** of the following intolerable and persistent (i.e. more than 2 weeks) symptoms for low or moderate-, and high-intensity statins:

1. Myalgia (muscle symptoms without CK elevations)
2. Myositis (muscle symptoms with CK elevations < 10 times upper limit of normal [ULN])

**-OR-**

- ii. Patient has a labeled contraindication to all statins as documented in medical records

**-OR-**

- iii. Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations > 10 times ULN

**-AND-**

- (3) One of the following:

(a) Submission of medical record (e.g., laboratory values) documenting **one** of the following LDL-C values while on maximally tolerated lipid lowering therapy within the last 120 days:

- i. LDL-C  $\geq$  100 mg/dL with ASCVD
- ii. LDL-C  $\geq$  130 mg/dL without ASCVD

**-OR-**

- (b) **Both** of the following:

i. Submission of medical record (e.g., laboratory values) documenting **one** of the following LDL-C values while on maximally tolerated lipid lowering therapy within the last 120 days:

1. LDL-C between 70 mg/dL and 99 mg/dL with ASCVD
2. LDL-C between 100 mg/dL and 129 mg/dL without ASCVD

**-AND-**

ii. Submission of medical record (e.g., chart notes, laboratory values) documenting **one** of the following [prescription claims history may be used in conjunction as documentation of medication use, dose, and duration]:

1. Patient has been receiving at least 12 consecutive weeks of ezetimibe (Zetia<sup>®</sup>) therapy as adjunct to maximally tolerated statin therapy

**-OR-**

2. Patient has a history of contraindication or intolerance to ezetimibe

**-AND-**

- (4) Used as an adjunct to a low-fat diet and exercise

**-AND-**

- (5) Prescribed by **one** of the following:

- (a) Cardiologist
- (b) Endocrinologist
- (c) Lipid specialist

**-AND-**

- (6) Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor

**-AND-**

- (7) **One** of the following as documented by prescription claims history or medical record:

- (a) History of failure after 12 consecutive weeks of Praluent at FDA maximum labeled dosing of 150 mg every 2 weeks

**-OR-**

- (b) History of intolerance to Praluent therapy

**-AND-**

- (8) Prescriber attests to the following: the information provided is true and

accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

\*Results of prior genetic testing can be submitted as confirmation of diagnosis of HeFH, however please note that UnitedHealthcare does not currently cover genetic testing for evidence of an LDL-receptor mutation, familial defective apo B-100 or a PCSK9 mutation.

**Authorization will be issued for 12 months**

2. **Reauthorization**

a. **Repatha** will be approved based on **all** of the following criteria:

- (1) Patient continues to receive statin at maximally tolerated dose (unless patient has documented inability to take statins)

**-AND-**

- (2) Patient is continuing a low-fat diet and exercise regimen

**-AND-**

- (3) Prescribed by **one** of the following:

- (a) Cardiologist  
(b) Endocrinologist  
(c) Lipid specialist

**-AND-**

- (4) Submission of medical records (e.g. chart notes, laboratory values) documenting LDL-C reduction while on Repatha therapy

**-AND-**

- (5) Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor

**-AND-**

- (6) Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

**Authorization will be issued for 12 months**

**B. Homozygous Familial Hypercholesterolemia**

**1. Initial Therapy**

a. **Repatha** will be approved based on **all** of the following criteria:

(1) Diagnosis of homozygous familial hypercholesterolemia (HoFH) as confirmed by submission of medical records (e.g., chart notes, laboratory values) documenting **both** of the following:\*

(a) **One** of the following:

- i. Pre-treatment LDL-C greater than 500 mg/dL
- ii. Treated LDL-C greater than 300 mg/dL

**-AND-**

(b) **One** of the following:

- i. Xanthoma before 10 years of age
- ii. Evidence of heterozygous familial hypercholesterolemia (HeFH) in both parents

**-AND-**

(2) Used as an adjunct to a low-fat diet and exercise

**-AND-**

(3) Patient is receiving other lipid-lowering therapy (e.g., statin, ezetimibe, LDL apheresis)

**-AND-**

(4) Prescribed by **one** of the following:

- (a) Cardiologist
- (b) Endocrinologist
- (c) Lipid specialist

**-AND-**

- (5) Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor

**-AND-**

- (6) Not used in combination with Juxtapid (lomitapide)

**-AND-**

- (7) Not used in combination with Kynamro (mipomersen)

**-AND-**

- (8) Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

\*Results of prior genetic testing can be submitted as confirmation of diagnosis of HeFH, however please note that UnitedHealthcare does not currently cover genetic testing for evidence of an LDL-receptor mutation, familial defective apo B-100 or a PCSK9 mutation.

**Authorization will be issued for 12 months.**

**2. Reauthorization**

a. **Repatha** will be approved based on **all** of the following criteria:

- (1) Patient is continuing a low-fat diet and exercise regimen

**-AND-**

- (2) Patient continues to receive other lipid-lowering therapy (e.g., statin, LDL apheresis)

**-AND-**

- (3) Submission of medical records (e.g. chart notes, laboratory values) documenting LDL-C reduction while on Repatha therapy



**-AND-**

(4) Prescribed by **one** of the following:

- (a) Cardiologist
- (b) Endocrinologist
- (c) Lipid specialist

**--AND-**

(5) Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor

**-AND-**

(6) Not used in combination with Juxtapid (lomitapide)

**-AND-**

(7) Not used in combination with Kynamro (mipomersen)

**-AND-**

(8) Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

**Authorization will be issued for 12 months.**

### 3. References:

1. Repatha [package insert]. Thousand Oaks, CA : Amgen Inc.; July 2016.
2. WHO Familial Hypercholesterolemia Consultation Group. Familial Hypercholesterolemia (FH): report of a second WHO consultation. Geneva: World Health Organization; 1999.
3. Scientific Steering Committee on behalf of the Simon Broome Register Group. Risk of fatal coronary heart disease in familial hypercholesterolaemia. *BMJ*. 1991;303:893-6.
4. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;63:2889-934.

Confidential and Proprietary, © 2017 UnitedHealthcare Services Inc.

5. Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*. 2015a; DOI: 10.1056/NEJMoa1410489 [Epub ahead of print].
6. The Lipid Research Clinics Coronary Primary Prevention Trial results. II. The relationship of reduction in incidence of coronary heart disease to cholesterol lowering. *JAMA*. 1984;251:365-74.
7. ATP III Final Report PDF. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. *Circulation*. 2002;106:3143-3421.
8. Per clinical drug consult with cardiologist. August 3, 2015.
9. Blom DJ, Hala T, Bolognese M, et al. A 52-week placebo-controlled trial of evolocumab in hyperlipidemia. *N Engl J Med*. 2014;370:1809-19.
10. Raal FJ, Santos RD. Homozygous familial hypercholesterolemia: current perspectives on diagnosis and treatment. *Atherosclerosis*. 2012;223:262-8.
11. Raal FJ, Honarpour N, Blom DJ, et al. Inhibition of PCSK9 with evolocumab in homozygous familial hypercholesterolaemia (TESLA Part B): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015;385:341-50.
12. Cuchel M, Bruckert E, Ginsberg HN, et al. Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. *Eur Heart J*. 2014;35:2146-57.
13. Lloyd-Jones D, Morris P, Ballantyne C, et al. 2016 ACC expert consensus decision pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk. *J Am Coll Cardiol*. 2016;68:92-125.
14. Austin MA, Hutter CM, Zimmern RL, Humphries SE. Genetic causes of monogenic heterozygous familial hypercholesterolemia: a HuGE prevalence review. *American journal of epidemiology*. 2004;160:407-420
15. Haase A, Goldberg AC. Identification of people with heterozygous familial hypercholesterolemia. *Current opinion in lipidology*. 2012;23:282-289.
16. Nordestgaard BG, Chapman MJ, Humphries SE, et al. Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease: consensus statement of the European Atherosclerosis Society. *European heart journal*. 2013;34:3478-3490a.
17. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American association of clinical endocrinologists and American college of endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. *Endocr Pract*. 2017; Suppl 2;23:1-87.

Program	Prior Authorization –Repatha (evolocumab)
<b>Change Control</b>	
Date	Change
5/2015	New program.

5/2015	Added examples of atherosclerotic cardiovascular disease.
9/2015	Revised clinical criteria to include combination use of high-intensity statin or documented intolerance to high-, moderate- and low intensity statin therapy to achieve the maximally tolerated statin therapy.
11/2015	Added step therapy requirement language for Primary Hyperlipidemia
1/2016	Removed continuation of therapy criterion.
8/2016	Add requirement of Praluent failure at maximum labeled dosing, Updated reference.
12/2016	Modified medical record criteria to include review of prescription claims history. Updated references.
3/2017	Changed initial authorization duration to 12 months for all indications
11/2017	Updated medical record requirement, modified criteria for HeFH diagnosis, modified previous statin requirement requiring failure, intolerance to high intensity and either moderate or low intensity statin. Modified target LDL values and ezetimibe trial requirement. Extended timeline for lipid panel submission to 120 days. Added physician attestation criterion. Updated references.